Author Index

Andrawes, F.
——, Greenhouse, S. and Draney, D.
Chemistry of acrylamide bromination for trace analysis by gas chromatography and gas chromatography–mass spectrometry 269
Barkley, R. M., see Pourreza, N. 165
Batten, C. F., see Chen, E. C. M. 121
Bertsch, W., see Mar, T. 277
Brill, J., see Mar, T. 277
Brooks, C. J. W.
—— and Cole, W. J.
Cyclic ferroceneboronates as derivatives for the gas chromatographic separation and characterisation of diols and related compounds 207
Bruner, F.
——, Crescentini, G., Mangani, F., Palma, P. and Xiang, M.
Performance and use of wall-coated open tubular fused-silica columns with liquid-modified graphitized carbon black 87
Burford, R. P., see haken, J. K. 295
Chamberlin, B. A., see Sweeley, C. C. 173
Chen, E. C. M.
——, Wentworth, W. E., Desai, E. and Batten, C. F.
Importance of kinetics and thermodynamics to the electron-capture detector 121
Cole, W. J., see Brooks, C. J. W. 207
Crescentini, G., see Bruner, F. 87
Crewe, R., see Mar, T. 277
Desai, E., see Chen, E. C. M. 121
Downing, G. V., see Wehner, T. A. 251
Draney, D., see Andrawes, F. 269
Eggstein, M., see Liebich, H. M. 291
Fischer, U., see Liebich, H. M. 291
Fletcher, D. J. C., see Mar, T. 277
Furton, K. G.
—— and Poole, C. F.
Thermodynamic characteristics of solute–solvent interactions in liquid organic salt solvents, studied by gas chromatography 47
Ghaoui, L.
Comparison of different adsorbents for isolation of polynuclear aromatic hydrocarbons from water 69
Greenhouse, S., see Andrawes, F. 269
Haken, J. K.
——, Vimalasiri, P. A. D. T. and Burford, R. P.
Chromatographic analysis of thermoplastic polyurethanes based on hydroxy-terminated polycaprolactones after alkali fusion 295
Hayashiya, K., see Yamaoka, R. 259
Holland, J. F., see Sweeley, C. C. 173
Horie, K., see Yamashita, K. 223
Horning, E.C.
——, Lertratanangkoon, K. and Horning, M. G.
Metabolism of bromobenzene. Analytical chemical and structural problems associated with studies of the metabolism of a model aromatic compound 321
——, see Horning, M. G. 303
Horning, M. G.
——, Sheng, L.-S., Nowlin, J. G., Lertratanangkoon, K. and Horning, E. C.
Analytical methods for the study of urinary thioether metabolites in the rat and guinea pig 303
——, see Horning, E. C. 321
Hung, L. B., see Parcher, J. F. 75
Hutte, R. S., see Pourreza, N. 165
Hyver, K. J.
—— and Phillips, R. J.
Considerations in enhancing resolution, speed, and sensitivity in capillary gas chromatography and gas chromatography–mass spectrometry 33
Imai, Y., see Tatsuhara, T. 183
Irie, T., see Ishibashi, M. 197
Ishibashi, M.,
——, Irie, T. and Miyazaki, H.
Diethylhydrogensilyl-cyclic diethylsilylene derivatives in the gas chromatography–mass spectrometry of hydroxylated steroids. IV. Hydrocortisone 197
——, see Yamashita, K. 223
Ito, T., see Mogi, M. 245
Jones, J. R., see Purnell, J. H. 99
Kersten, B. R.
—— and Poole, C. F.
Influence of concurrent retention mechanisms on the determination of stationary phase selectivity in gas chromatography 1
Kurata, Y., see Mogi, M. 245
Lertratanangkoon, K., see Horning, E. C. 321
——, see Horning, M. G. 303
Liebich, H. M.
——, Risler, T., Fischer, U., Rapp, K., Tetschner, B. and Eggstein, M.
Gas chromatographic determination of hippuric acid and a furancarboxylic acid in plasma of patients with kidney diseases 291
Mangani, F., see Bruner, F. 87

Mar, T.
——, Brill, J., Bertsch, W., Fletcher, D. J. C. and Crewe, R.
Investigation of cuticular hydrocarbons from selected honeybee populations by gas chromatography with pattern recognition 277
Matsuda, Y., see Tatsuhara, T. 183
Matsuki, Y., see Mogi, M. 245
Miyazaki, H., see Ishibashi, M. 197
——, see Yamashita, K. 223
Mogi, M.
——, Ito, T., Matsuki, Y., Kurata, Y. and Nambara, T.
Determination of isofloxythepin in biological fluids by gas chromatography–mass spectrometry 245
Montzka, S. A., see Pourreza, N. 165
Muro, H., see Tatsuhara, T. 183
Nambara, T., see Mogi, M. 245
Niemelä, K.
Oxidative and non-oxidative alkali-catalysed degradation of L-ascorbic acid 235
Nowlin, J. G., see Horning, M. G. 303
Palma, P., see Bruner, F. 87
Parcher, J. F.
—— and Hung, L. B.
Cooperative adsorption effects in gas–liquid–solid chromatography 75
Phillips, R. J., see Hyver, K. J. 33
Poole, C. F., see Furton, K. G. 47
——, see Kersten, B. R. 1
Pourreza, N.
——, Montzka, S. A., Barkley, R. M., Sievers, R. E. and Hutte, R. S.
Chemistry occurring in redox chemiluminescence detectors for chromatography 165
Purnell, J. H.
——, Jones, J. R. and Wattan, M. H.
Theory and practice of coupled gas chromatographic columns 99
Ramsey, R. S.
—— and Todd, R. A.
Pulsed-modulated helium-ionization detection 139
Rapp, K., see Liebich, H. M. 291
Risler, T., see Liebich, H. M. 291
Sheng, L.-S., see Horning, M. G. 303
Sievers, R. E., see Pourreza, N. 165
Simmonds, P. G.
Electron-capture detector with a photoemissive electron source 149
Sweeley, C. C.
——, Holland, J. F., Towson, D. S. and Chamberlin, B. A.
Interactive and multi-sensory analysis of complex mixtures by an automated gas chromatography system 173
Tatsuhara, T.
——, Muro, H., Matsuda, Y. and Imai, Y.
Determination of valproic acid and its metabolites by gas chromatography–mass spectrometry with selected ion monitoring 183
Tetschner, B., see Liebich, H. M. 291
Todd, R. A., see Ramsey, R. S. 139
Tokoro, M., see Yamaoka, R. 259
Towson, D. S., see Sweeley, C. C. 173
VandenHeuvel, W. J. A., see Wehner, T. A. 251
Vimalasiri, P. A. D. T., see Haken, J. K. 295
Walker, R., see Wehner, T. A. 251
Watabe, K.
Resolution of α-halogenocarboxylic acid amides on chiral N-lauroyl-α-(1-naphtyl) ethylamine. Effect of the nature of the amide moiety of the solute 111
Watanabe, K., see Yamashita, K. 223
Wattan, M. H., see Purnell, J. H. 99
Wehner, T. A.
——, Wood, Jr., J. S., Walker, R., Downing, G. V. and VandenHeuvel, W. J. A.
Confirmation of clorsulon residues in cattle kidney by capillary gas chromatography–negative-ion chemical-ionization mass spectrometry 251
Wentworth, W. E., see Chen, E. C. M. 121
Wood, Jr., J. S., see Wehner, T. A. 251
Yamamoto, S., see Yamashita, K. 223
Yamaoka, R.
——, Tokoro, M. and Hayashiya, K.
Determination of geometric configuration in minute amounts of highly unsaturated termite trail pheromone by capillary gas chromatography in combination with mass spectrometry and Fourier-transform infrared spectroscopy 259
Yamashita, K.
——, Watanabe, K., Ishibashi, M., Miyazaki, H., Yokota, K., Horie, K. and Yamamoto, S.
Gas chromatography–mass spectrometry of some prostanoids with new derivatizing agents. Application to the analysis of 13,14-dihydro-15-ketoprostaglandin $F_{2\alpha}$ 223
Yokota, K., see Yamashita, K. 223
Xiang, M., see Bruner, F. 87